

Journal Pre-proof

A transdiagnostic meta-analysis of physical and social anhedonia in major depressive disorder and schizophrenia spectrum disorders

Arti Gandhi , Jasmine Mote , Daniel Fulford

PII: S0165-1781(21)00673-9
DOI: <https://doi.org/10.1016/j.psychres.2021.114379>
Reference: PSY 114379



To appear in: *Psychiatry Research*

Received date: 7 June 2021
Revised date: 9 December 2021
Accepted date: 29 December 2021

Please cite this article as: Arti Gandhi , Jasmine Mote , Daniel Fulford , A transdiagnostic meta-analysis of physical and social anhedonia in major depressive disorder and schizophrenia spectrum disorders, *Psychiatry Research* (2022), doi: <https://doi.org/10.1016/j.psychres.2021.114379>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Published by Elsevier B.V.

A transdiagnostic meta-analysis of physical and social anhedonia in major depressive disorder and schizophrenia spectrum disorders

Arti Gandhi¹, Jasmine Mote², Daniel Fulford^{1,3}

¹Sargent College of Health and Rehabilitation Sciences, Boston University
635 Commonwealth Ave,
Boston MA, 02215

² Department of Occupational Therapy, Tufts University
574 Boston Avenue
School of Arts and Sciences
Tufts University
Medford, MA 02155

³Department of Psychological & Brain Sciences, Boston University
900 Commonwealth Ave,
Boston MA, 02215

Highlights

- Anhedonia is elevated in Schizophrenia (SZ) and Major Depression
- People with SZ experience elevated social anhedonia compared to physical anhedonia
- Clinical severity moderates physical but not social anhedonia in Major Depression

Abstract

Introduction: Anhedonia is a transdiagnostic construct conceptualized as physical or social, however, the extent to which these subtypes differ across psychotic and mood pathology remains poorly understood. We aimed to quantify the severity of physical and social anhedonia across Major Depressive Disorder (MDD) and Schizophrenia Spectrum Disorder (SSDs).

Methods: We conducted meta-analyses of the Chapman Physical and Social Anhedonia Scales (PAS;SAS). We reviewed data from participants with MDD, and SSDs separately.

Results: Our first meta-analysis (n= 8 studies, 409 participants) with MDD revealed elevated SAS and PAS in MDD compared to controls. Within-group differences were not significant. Depressive symptom severity moderated the between-group effect of PAS. Our second meta-analysis (n= 44 studies, 3352 participants) revealed elevated SAS and PAS in SSDs compared to controls. We detected a moderate difference between the SAS and PAS within the SSD group. Age moderated within-group differences of SAS and PAS.

Discussion: People with SSD or MDD experience elevated SAS and PAS compared to controls. People with SSDs endorse greater challenges experiencing social rewards relative to physical rewards. People with MDD experience social and physical rewards similarly. The moderating role of depressive symptoms in MDD suggests that physical anhedonia is more state-like than social anhedonia.

Introduction

Anhedonia is characterized by a reduced experience of pleasure, at least partly reflected by impaired reward processing (Zhang et al., 2016), that can limit goal-directed activity and associated psychosocial functioning (Barch & Dowd, 2010; Kring & Barch, 2014). As a Diagnostic and Statistical Manual (DSM-5) Criterion A symptom of Major Depressive Disorder, anhedonia is central to the diagnosis of depression. It is also central to the diagnosis of Schizophrenia Spectrum Disorders (i.e., Schizophrenia and Schizoaffective Disorder) as one of five core negative symptoms (APA, 2013). Furthermore, anhedonia is a risk factor for the onset of both SSDs (Gooding et al., 2005; Kwapil et al., 1997; Velthrost et al., 2009) and MDD (Bress et al., 2013; Morgan et al., 2013). Anhedonia is thus a key transdiagnostic symptom and

common intervention target (Barkus & Badcock, 2019; Bedwell, Gooding, Chan, & Trachick, 2014).

Early conceptualizations considered the experience of anhedonia to be similar across SSDs and MDD. However, recent innovations in the assessment of emotion experience have painted a more complex and multifaceted picture. Specifically, assessments of “in-the-moment,” or consummatory pleasure experiences (i.e., ‘liking’), on the whole do not find differences in pleasant emotion experience between people with versus without SSDs. The assessment of in-situ responses to pleasant smells, images, social interactions, or other stimuli, using event-based ecological momentary assessment (EMA) and laboratory tasks, shows that people with SSDs demonstrate the capacity to experience pleasure (Blanchard, Mueser, & Bellack, 1998; Burbridge & Barch, 2007; Cohen et al., 2005; Horan & Blanchard, 2003, Horan et al., 2006). However, trait-based assessments (i.e., those that ask people to report a gestalt view of their emotional experiences) continue to show general pleasure deficits across daily life in people with SSDs compared to those without (Cho, Gonzalez, Lavaysse, Pence, Fulford, & Gard, 2017; Culbreth et al., 2016; Edwards, et al., 2015, Gerritsen, 2015). Other self-report measures and lab-based studies consistently demonstrate deficits in anticipatory pleasure (i.e., looking forward to future events and/or feeling pleasure in anticipation of those events; also known as ‘wanting’) among people with SSDs (Gard, Kring, Gard, Horan, Green, 2007). Taken together, people with SSDs appear to retain the capacity to experience and report the same amount of consummatory pleasant emotions compared to people without SSDs, yet show deficits when reporting trait-based and anticipatory pleasure experiences. Thus, rather than a general hedonic deficit, low positive affect typical in SSDs may instead reflect cognitive alterations in the way hedonic experiences are stored, represented, maintained, and ultimately reported (Barch & Dowd, 2019; Kring & Barch, 2014).

There is evidence that anhedonia in the context of MDD, however, may cut across state- (consummatory and anticipatory) and trait-based pleasure experiences. Regardless of the

method of assessment, people with MDD report reduced pleasure across multiple contexts (e.g., physical, interpersonal, monetary rewards; Barch, Pagliaccio, and Luking, 2015; Bylsma et al., 2008). Furthermore, there is less evidence for a contribution of impaired cognitive control systems to anhedonia in MDD, unlike in SSDs (Barch et al., 2015). These findings suggest that, unlike in SSDs, anhedonia in MDD may represent a reduced capacity to experience pleasure, more consistent with a generalized hedonic deficit.

Other innovations in the assessment of anhedonia reflect the different *types* of stimuli used to evoke emotion. The vast majority of studies assess anhedonia in the context of either physical (e.g., pleasant smells, images, sounds) or social (e.g., dynamic interpersonal interactions) stimuli. Lab-based tasks using static, sensory stimuli demonstrate consistent and robust findings of physical anhedonia among those with MDD (Barch et al., 2015; Bylsma et al., 2008) but intact affective responding among those with SSDs (Barch et al., 2015). Similarly, EMA studies assessing anhedonia in the context of real-life social interactions suggest that individuals with SSDs may demonstrate normative social pleasure in daily life (Mote & Fulford, 2020), while depressed individuals report social pleasure deficits in these contexts (Silk et al., 2011).

To date, few studies have directly compared physical and social anhedonia within or across groups. It may be that social and physical anhedonia differentially impact meaningful outcomes that could be targeted by different treatment approaches. Some recent research has found that social anhedonia, for example, is associated with functional deficits, illness severity, and other aversive outcomes such as loneliness, suicidality, and poor quality of life across both SSDs and MDD (Ritsner, Ratner, Mendyk, Gooding, 2018; Sagud et al., 2020). Furthermore, preliminary findings from neuroimaging and behavioral studies indicate blunted responses to social relative to nonsocial rewards across SSDs (Bjorkquist & Herbener, 2012; Catalano, Heerey, and Gold, 2018; Lee, Jimenez, Reavis, Horan, Wynn, & Green, 2018). Although such findings are inconsistent with previous EMA studies indicating increased positive affect in social

contexts within SSDs, they suggest that at on a neural and behavioral level, anhedonia may manifest differently across social and non-social stimuli (Fulford et al., 2018).

The most commonly used measures of trait-based social and physical anhedonia are the Chapman Anhedonia Scales (Chapman et al., 1976; Eckblad et al., 1982). They were the original scales designed to examine physical (The Physical Anhedonia Scale [PAS]) and social (The Social Anhedonia Scale [SAS]) anhedonia as separate constructs. Together, the scales reflect a quick, viable means of detecting trait-level anhedonia across contexts, including over time (e.g., test-retest reliability over 6-month intervals) and between cultures (e.g., North American and Chinese college students; Chan et al., 2014). Early studies in clinical populations showed that people with SSDs reported both physical and social anhedonia stably over time (Blanchard et al., 1998; Herbener & Harrow, 2002), while those with MDD reported anhedonia more episodically, covarying with clinical severity (Blanchard et al., 1998; Shankman et al., 2010).

Other studies highlight the potential clinical utility of the scales. Among those with SSDs, the Chapman Scales are negatively correlated with positive affect, social functioning, and pre-morbid functioning (Blanchard et al., 1998). Within MDD, higher PAS scores are associated with poorer social functioning and increased rates of rehospitalization over time (Shankman et al., 2010). Both scales are also positively correlated with lifetime suicide attempts and suicidal ideation in MDD (Sagud et al., 2020). Furthermore, psychometric assessments of the Chapman Scales found that approximately 24% of healthy individuals with elevated SAS scores endorsed an SSD diagnosis at the 10-year follow-up, compared to only 1% of those with elevated PAS scores (Chapman, Chapman, Kwapil, Eckblad, & Zinser, 1994; Kwapil, Miller, Zinser, Chapman, and Chapman, 1997; Kwapil, 1998). Thus, trait-like anhedonia, as assessed by the Chapman Scales, provides useful clinical information, such as the severity of psychopathology and related need for intervention. Furthermore, compared to the PAS, the SAS may provide greater utility as a predictor of the development of severe psychopathology over time among healthy individuals.

Despite the utility and ubiquity of the use of the Chapman Scales (as well as the differentiation between social and physical anhedonia broadly) in psychopathology research, there is limited evidence that trait-based physical and social anhedonia vary in their relative presentation in mood and psychotic disorders. Quantification of such differences may have important implications for how trait-based anhedonia contributes to psychosocial impairment, within and across disorders. For example, if trait-based social anhedonia is more pronounced in SSDs vs. controls, relative to MDD vs. controls, such impairment could reflect different biopsychosocial mechanisms that could be targeted in treatments. In other words, if social or physical anhedonia is more primary in some diagnoses, and more generalized in others, it may clarify the granularity of anhedonia, informing more precise treatment approaches. As emotion experience assessments continue to advance in differentiating between consummatory, anticipatory, and other trait-based experiences, it remains vital to understand the utility of continuing to view trait-based social and physical anhedonia as separable constructs among two of the most debilitating mental health concerns.

In the current-meta-analysis we quantified the within- and between-group (i.e., clinical vs. controls) differences in trait-like, self-reported physical and social anhedonia across MDD and SSDs, specifically, those with Schizophrenia and Schizoaffective disorder. First, we examined the magnitude of difference between each diagnostic group and controls on the Chapman Scales. Next, we quantified the discrepancies between physical and social anhedonia *within* each clinical group by examining the magnitude of difference between means scores on the Chapman Scales. We further aimed to investigate the degree to which clinical symptoms, including depression and negative symptoms, moderated between-group differences in physical and social anhedonia. Although anhedonia is considered a negative symptom, we examined interview-rated negative symptom severity as a moderator of Chapman Scale scores to identify the degree to which the scales capture variance in gold-standard clinical measures. Finally, we

examined additional conditions and characteristics as moderators of such differences, including age, sex, race, and mood episode and medication status.

Methods

This review was registered on July 2020 on the Open Science Framework (OSF) with a PRISMA guided protocol: <https://osf.io/pmk2c/>.

Search Strategy

We conducted two separate searches of articles published between January 1976 (the year the Chapman Anhedonia scales were published) and April 2020 using the PubMed and PsycINFO databases. We first searched for articles assessing anhedonia in MDD using the following search terms: “major depressive disorder” or “MDD” or “major depression” and “social” or “physical” and “anhedonia” or “pleasure”. Next, we searched for articles assessing anhedonia and schizophrenia using similar search terms: “schizo*” or “psychosis” and “social” or “physical” and “anhedonia” or “pleasure”. We also used professional listservs to identify any unpublished data to incorporate in the analyses. We used the following inclusion criteria: (1) samples consisting of SSDs and/or MDD, (2) adults (ages 18 or older), (3) empirical study published in a peer-reviewed journal, (4) in the English language or translated, (5) using the Chapman Anhedonia Scales, and (6) the revised version of the SAS. We restricted our search to the revised SAS as the original SAS was found to include items tapping into social anxiety, rather than social anhedonia (Eckblad, Chapman, and Chapman, 1982). We also used the following exclusion criteria: (1) systematic reviews or (2) case studies; (3) studies with participants with co-occurring substance use disorders, (4) Schizotypal Personality Disorder (5) at clinical high-risk for psychosis; (6) studies with no control group. Co-occurring substance use was excluded from the analysis as substance use may artificially impact ventral striatal and other motivation and reward pathways relevant for anhedonia. Furthermore, we only included studies using Chapman Scales with a factor structure that is congruent with the English version.

Data Extraction & Coding

We uploaded search results into the Covidence systematic review website (<https://www.covidence.org>). Two separate reviewers—the first-author and a trained research assistant—reviewed the abstract and full-text for both searches. Any discrepancies were discussed by both reviewers until a consensus was reached. Reviewers extracted means, standard deviations, and sample sizes for each scale across both clinical and control groups. When means or standard deviations were not available, authors were contacted for either raw data or summary scores. Reviewers extracted the following covariates: (1) mean age, (2) percentage of white participants, (3) percentage of men, (4) percentage of atypical antipsychotics prescribed to SSD samples, (5) percentage of antidepressants prescribed to MDD samples, (6) mean and standard deviation of depressive symptom severity, (7) MDE vs. MDD status, and (8) mean and standard deviation for negative symptoms within SSD samples only. To allow for pooling across various scales, negative symptom and depression scores were transformed to proportions by dividing the mean values by the total points possible on each measure, and then multiplying by 100. Authors were contacted when data were not available.

We conducted the searches in April 2020 (See figure 1 for flowchart of study selection and exclusion process). The SSD search yielded 495 results, of which 116 duplicates were removed, leaving 379 for review. After abstract screening and full-text review, a total of 44 studies with 44 individual effects were included for review. See Tables 1.a-b for details of study characteristics. The MDD search yielded 283 results, of which 36 duplicates were removed, leaving 247 for review. See Figure 2 for flowchart of study selection and exclusion process. Following abstract screening and full-text review, a total of 8 studies were included for review. See Table 2 for details of study characteristics.

Finally, we used Web of Science to search for studies that cited the Chapman scales. We found 308 studies including MDD groups, and 612 studies including SSD groups. After comparing these results from our previous PsycInfo and PubMed searches, and eliminating duplicates, we identified 117 SSD articles that were not previously screened, and 25 MDD

articles that were not previously screened. After screening the title and abstract, none of these additional articles met inclusion criteria.

Statistical Analysis

We calculated Hedge's g for differences between clinical and control groups across PAS and SAS scales by dividing the difference of clinical and control means by the pooled standard deviation. To assess the magnitude of difference between PAS and SAS within each clinical group, we calculated Cohen's d for repeated measures using the following formula (Cohen's d_{rm} ; Cohen, 1988; Lakens, 2013):

$$\text{Cohen's } d_{rm} = \frac{M_{diff}}{\sqrt{SD1^2 + SD2^2 - 2 \times r \times SD1 \times SD2}} \times \sqrt{2(1 - r)}$$

The numerator represents the difference of the standardized PAS and SAS means, while the denominator represents the standard deviation of differences. We used $r = 0.40$ to represent the correlation between PAS and SAS measures, as this correlation was established in several prior studies (e.g., Chapman et al., 1995; Fonesca-Pedrero et al., 2008; Kwapil, 2008). We then converted these values to Hedge's g to account for small sample bias. We conducted all analyses using the *dmetar* package (Harrer, Cuijpers, Furukawa, and Ebert, 2019) on R software (R Core Team, 2021). To determine model heterogeneity, we used Q and I^2 (Higgins et al., 2003). I^2 values 25% or below represent low heterogeneity, while values above 25% represent moderate to high heterogeneity (Higgins et al., 2003). When I^2 was greater than 25%, we ran random effects models, and for I^2 values below 25%, we ran fixed effects models.

For each effect, we detected outliers by searching for studies with confidence intervals (CIs) that did not overlap with the CI of the pooled effect. In other words, we defined a study as an outlier if 1) the upper bound of the 95% CI was lower than the lower bound of the pooled effect CI, or 2) if the lower bound of the 95% CI was higher than the upper bound of the pooled

effect CI (Viechtbauer & Cheung, 2010). We report effects both with and without outliers included in overall pooling.

We examined the influence of negative symptoms, depressive symptoms, age, sex, race and medication status on between-group and within-group differences across both scales, by conducting mixed-effects linear regression models. For the MDD studies, we did not assess the influence of remitted MDD vs. current MDD, race, or antidepressants as a moderator as there was not a sufficient number of studies providing the necessary data. Finally, we examined publication bias through funnel plots and Egger's tests for each pooled effect. When interpreting funnel plots, we examined the extent to which the studies evenly scattered on either side of the overall effect. An uneven scattering of studies may suggest the presence of publication bias.

Results

Schizophrenia Spectrum Disorders

Analyses of PAS between SSD and controls yielded a large effect ($g = 0.89$; 95% CI: 0.78-1.00; $t = 16.12$; $p < 0.0001$), indicating that individuals with SSD self-reported greater physical anhedonia than controls. This effect increased to 0.97 (95% CI: 0.88 – 1.10; $t = 22.94$; $p < 0.0001$) after removing four outliers (Baslet et al., 2009; Burbridge et al., 2007; Makowski et al., 2016; Zou et al., 2018). Group differences on SAS yielded a similarly large effect ($g = 0.93$; 95% CI: 0.67 – 1.18; $t = 7.32$, $p < 0.0001$), demonstrating that individuals with SSD self-reported greater social anhedonia than controls. We detected one extreme outlier (Umesh et al., 2018), which yielded an effect size of 8.7. Upon further investigation of this study, we found that standard deviations for both controls and SSDs were extremely low (i.e., 0.97 and 1.40, respectively). Nevertheless, after removing this outlier, the group difference in SAS remained large ($g = 0.84$; 95% CI: 0.76 – .92; $t = 20.80$; $p < 0.0001$). The Q statistic was significant for each effect suggesting heterogeneity beyond sampling error. I^2 values indicate that less than 60% of the observed variance in each effect was due to sampling error (see Table 3). Mixed effects meta-regression models did not indicate any significant moderators across either effect.

Within-group differences between PAS and SAS in the SSD studies were pooled through a fixed effects model. This model yielded a moderate effect ($g = -0.50$; 95% CI: $-0.83 - -0.17$; $z = -3.00$; $p < 0.0001$). This suggests that SAS scores were significantly higher than PAS scores across SSD studies. The Q statistic was not significant, and $I^2 = 0.0\%$, suggesting no heterogeneity in the effects beyond sampling error (see Table 3). The effect size increased to -0.58 (95% CI: $-0.92 - -0.25$; $z = -3.45$; $p < 0.001$) after removing one outlier (Makowski et al., 2016). Mixed effects models indicated a significant effect of age on within-group differences between SAS and PAS, such that older individuals demonstrated smaller discrepancies between physical and social anhedonia compared to younger individuals. The remaining moderators were not significant. Of note, the average standardized score (0-100) for negative symptoms was 36.52.

Visual inspection of funnel plots and Egger's test revealed asymmetry and publication bias for the between-group effect of social anhedonia ($t(44) = 2.60$, $p = 0.01$), and the within-group difference between PAS and SAS ($t(41) = -2.09$, $p = 0.04$). For the between group effect, asymmetry appeared to be primarily driven by the Umesh et al. (2018) study. For the within-group effect, asymmetry appeared to be driven by the Makowski et al. (2016) study. We did not find asymmetry or publication bias for the between-group effect of physical anhedonia ($t(41) = 1.42$, $p = 0.16$) (see Figures 9-11).

Major Depressive Disorder

Group differences in PAS scores between MDD and controls yielded a large effect of 1.18 (95% CI: $0.91 - 1.44$; $z = 8.68$; $p < 0.0001$), indicating that individuals with MDD report greater physical anhedonia than controls. There was also a large aggregated group difference in SAS scores ($g = 1.32$; 95% CI: $1.04 - 1.61$; $z = 9.04$; $p < 0.0001$), indicating greater self-reported social anhedonia in MDD compared to controls. Finally, the mean difference between PAS and SAS within MDD was not significant ($g = -0.72$; 95% CI: $-2.9 - 1.47$; $z = -0.64$; $p < 0.52$), indicating that individuals with MDD report similar social and physical anhedonia. No

outliers were detected for any of these effects. Q statistics suggested no heterogeneity beyond sampling error. I^2 values indicated that less than 40% of the observed variance in each effect was due to sampling error (see Table 4).

Mixed effects models indicated significant effect of depressive symptoms, such that greater symptoms were associated with a larger difference in PAS scores between those with MDD and controls. The remaining moderators of the MDD-Control group differences and within-group MDD differences were not significant. Funnel plots and Egger's tests did not indicate asymmetry or publication bias for any effect. However, because the sample size is small (8 studies), the Egger's test may lack statistical power to detect significance (see Figures 12-14).

Although underpowered (five studies each), we conducted two final analyses of studies that compared the Chapman Scales between MDD and SSDs directly (Berlin, Givry-Steiner, Lecrubier & Puech, 1998; Chuang et al., 2014; Olsen, Bjorkquist, Bodapati, Shankman & Herbener, 2015; Pelizza & Ferarri, 2009; Wang et al., 2020) Group differences indicate nonsignificantly elevated scores among those with MDD compared to those with SSDs on both PAS ($g=0.15$, $p=.17$) and SAS ($g=0.03$, $p=.78$). Across both meta-analyses, two studies showed negligible differences between the groups (Berlin et al 1998; Wang et al., 2020) two other studies showed moderate negative differences (i.e., SSDs elevated relative to MDD; Chuang et al., 2014; Olsen et al., 2015), and one study showed a large positive effect (i.e., MDD elevated relative to SSDs, Pelizza & Ferarri, 2009). The overall positive SMDs were likely driven by the one moderate positive effect, suggesting that the analysis is too underpowered to yield a meaningful interpretation. For more information, please see our OSF page <https://osf.io/m7fhw/>

Discussion

We conducted a meta-analytic review of trait-level, self-reported physical and social anhedonia across MDD and SSDs. We quantified the magnitude of physical and social anhedonia between controls and clinical groups, as well as between facets of anhedonia within

groups, and determined the extent to which clinical and demographic characteristics moderated effects. As predicted, people with SSDs and MDD reported significantly higher trait-level physical and social anhedonia than controls, reflected in large between-group effect sizes. These findings demonstrate a high degree of self-reported physical and social anhedonia across the published literature in clinical populations characterized by deficits in the experience of trait-based pleasure.

Elevated Chapman Scale scores across MDD and SSDs indicate that although anhedonia severity in MDD may be more state-like than in SSDs, both groups experience at least some degree of anhedonia in stable manner. In MDD, anhedonia may be more pronounced during periods of low mood, while in SSDs anhedonia may be elevated regardless of mood state; this suggests that although both groups experience elevated anhedonia, it may be of a different nature. Our results further demonstrated that trait-based social anhedonia may be more severe than physical anhedonia within SSDs, but similar across the social and physical domains within MDD. This suggests that people with SSDs endorse greater challenges experiencing pleasure from social rewards as opposed to non-social rewards, and thus provides preliminary support for treating trait-based physical and social anhedonia as distinct constructs. Elevated social anhedonia relative to physical anhedonia in SSDs aligns with previous lab-based elicitation tasks, demonstrating blunted in-the moment hedonic responses to social relative to nonsocial stimuli in SSD (Bjorkquist & Herbener, 2012; Catalano et al., 2018; Lee, Jimenez, Reavis, Horan, Wynn, & Green, 2018). In contrast, these studies are inconsistent with EMA reported findings of elevated positive affect in social contexts (Mote & Fulford, 2020). Future studies should re-examine potential differences in social and nonsocial (i.e., physical or monetary) stimuli across psychopathology to determine whether elevated social anhedonia (relative to nonsocial anhedonia) is specific to psychotic illnesses or is a transdiagnostic phenomenon.

Several challenges limit how we interpret the above findings. Within SSDs, elevated SAS relative to PAS may be driven by limited opportunities for social contact, rather than limited social hedonic capacity. Items such as “I don’t really feel very close to my friends” and “I like to make long distance phone calls to friends and relatives” often depend on the presence of recent interactions and ongoing relationships, which are consistently low in people with SSDs (Gayer-Anderson & Morgan, 2013) and MDD (Kupferberg, 2016). Indeed, reduced social network size, and a lower likelihood of being in romantic partnerships (Gayer-Anderson & Morgan, 2013; Kupferberg, 2016) is common in both these populations. Elevated trait-level social anhedonia may reflect impoverished social environments rather than dampened capacity to experience social pleasure. This may further clarify why EMA studies, which probe participants to consider in-the-moment experiences, indicate intact social pleasure (Mote & Fulford, 2020). A critical future direction is to examine the extent to which frequency of social contact is related to trait-based reports of social anhedonia.

In this study, limited social opportunities may also explain the relatively higher social anhedonia in the SSDs group but not in the MDD group. Although typically reduced in both groups, limited social opportunities may be even more pronounced in those with SSDs compared to those with MDD, since SSDs symptoms are more chronic and enduring. It is also important to note that many items in the SAS tap into asociality (i.e., another negative symptom describing the reduction of social initiation due to decreased interest in forming close relationships with others), which is a related, but distinct construct that typically captures attitudes and behavior arising from anhedonia, rather than the experience of anhedonia itself. Similarly—in-line with burgeoning research—elevated SAS may arise from challenges associated poor social cognition, and challenges with translating the desire for social affiliation to actual social behavior (Barch et al., 2015; Fulford et al., 2018).

Furthermore, consistent with most trait-level self-report scales, the Chapman Scales require individuals to reflect on past experiences that, due to retrospective recall biases, may

lead people to report experiencing less pleasure than they actually experienced when in-the-moment. Previous findings suggest that self-reported or clinician-rated anhedonia is more reflective of anticipatory than consummatory deficits, as individuals engage in similar memory recall when anticipating future events (Kring & Elis, 2013). Anticipatory pleasure deficits may also explain why findings of elevated trait-based anhedonia in SSD vs. controls, along with previous self-report findings, are inconsistent with laboratory-based tasks assessing in-the-moment responses among people with SSDs. Given that negative memory biases have been observed in both groups, future work should focus on comparing anhedonia across different methods to more directly examine the impact of recall bias on hedonic deficits.

Regarding our second aim, we were surprised to find that interview-rated negative symptom severity did not moderate group differences in physical or social anhedonia between those with SSDs and controls. Though unexpected, this finding is consistent with those of Visser et al. (2020), who found that negative symptoms did not moderate self-reported trait anhedonia in people with SSDs. On one hand, the lack of moderation from negative symptoms may be due to an insufficient number of SSD studies included in this review, leading to a lack of power in the analysis. On the other hand, these results may indicate that negative symptoms do not influence the variance of self-reported trait anhedonia. Unlike clinical interviews, self-report measures only capture the experiential component of negative symptoms (i.e., anhedonia, avolition, asociality) but not the expressive component (i.e., affective flattening, alogia). Thus, it is possible we were not able to detect moderation because the Chapman Scales do not capture the full scope of negative symptoms. Furthermore, responses on trait-based assessments may be more directly impacted by cognitive difficulties and negative memory biases than are those elicited by clinical interviews of negative symptoms. Finally, the lack of moderation may also reflect a lack of coherence between clinical interviews (i.e., observer-rated assessment) and self-reports, as this has been detected in previous studies (Durand, 2015; Gould et al., 2015).

Furthermore, we found that depressive symptom severity moderated group differences in physical anhedonia, but not social anhedonia, in people with MDD. These findings suggest that physical anhedonia, relative to social anhedonia, is more likely to covary with depressive symptoms, and therefore worsen as a function of depression. Further, these findings suggest that in regards to assessment and treatment, physical anhedonia may be a stronger marker for the presence of a depressive episode, while social anhedonia may persist as a trait-like deficit regardless of depressive symptom severity. Overall, it is possible that physical anhedonia covaries more strongly with the experience of depression than does social anhedonia. Nonetheless, given the relatively low magnitude of these effects, findings should be further examined in future studies.

Sample characteristics (i.e., age, sex, race, medication status) did not moderate within- or between-group differences in social or physical anhedonia in MDD. Within SSDs, age was a source of heterogeneity for the within-group discrepancy between physical and social anhedonia, such that older age was associated with smaller differences between the two forms of anhedonia. This finding is consistent with a recent epidemiological investigation which found lower social anhedonia in healthy older adults compared to middle-aged adults (Dodell-Feder & Germine, 2018). Accordingly, smaller differences between the two subtypes may reflect social anhedonia following a normative decline within SSD, with physical anhedonia persisting at relatively the same level. Indeed, if self-reported social anhedonia reflects limited social opportunities, as discussed above, it may be that higher social anhedonia in older age reflects reduced social interactions over time.

Finally, it is important to note potential for publication bias based on exclusion of two influential studies in our analyses. Umesh et al. (2018) was the only study included in this review that was conducted in India, while participants in Makowski et al. (2016) were all male. It is possible these study details contributed to extreme effect sizes. Nevertheless, the results did

not drastically change with the exclusion of these outliers, and therefore does not skew our interpretation of the results.

Limitations & Conclusions

Several limitations of this meta-analysis must be considered. First, the small number of MDD studies raises concerns for these analyses lacking statistical power to detect differences. It is unclear whether the null findings for the moderators and the SAS/PAS within-group differences reflect a lack of statistical power, or true null effects. Future studies should directly compare social and physical anhedonia in MDD to further clarify whether anhedonia generalizes across domains or is specific to the social domain. Second, we could only include a limited number of moderators, due to limited reporting of data. Consequently, it remains unclear how racialized experiences, active vs. recovered MDD status, or antidepressants may impact the effects of anhedonia in MDD. Third, the large degree of heterogeneity within the SSD studies limits the precision of the overall effects and challenges the generalizability of these findings. Fourth, due to the limited availability of data, we were unable to directly compare the effects of anhedonia between MDD and SSD. Finally, MDD findings may not generalize to those in remission, as the majority of our sample consisted of those experiencing a MDE at time of data collection.

In summary, this meta-analysis found that individuals with SSD and MDD reported greater trait social and physical anhedonia compared to controls. Importantly, only the SSD group demonstrated significant differences between social and physical anhedonia, while differences were not significant in MDD. Within MDD, depressive symptoms appear more strongly associated with physical anhedonia compared social anhedonia, and thus may be more predictive of a current episode. This is the first study to examine the relative difference between social and physical anhedonia using one of the most widely used scales in psychopathology literature. Our results add to the literature by suggesting that the extent to which trait-level, self-reported anhedonia is primarily physical, social, or generalized in nature may differ across

clinical groups. Specifically, for people with SSDs, the experience of anhedonia may be more social in nature, while for people with MDD, the experience of anhedonia is more generalized. These results provide evidence for differing phenomenology and etiology of anhedonia across mood and psychotic disorders, and may inform more refined treatment targets.

Key Words

1. Anhedonia
2. Schizophrenia
3. Depression

Declarations of Interest: none

Author statement

Arti Gandhi: Conceptualization; methodology; formal analysis; Writing-Original & Review; Visualization

Jasmine Mote: Conceptualization; Writing-Original & Review

Daniel Fulford: Conceptualization; Writing- Original & Review, Methodology, Supervision

References

Alan Edward Ceaser, & Deanna M Barch. (2016). Striatal activity is associated with deficits of cognitive control and aberrant salience for patients with schizophrenia. *Frontiers in Human Neuroscience*, 9(2016), 687.

American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Arlington, VA: Author.

- Assaad, L., & Lemay, E. P. (2017). Social anhedonia and romantic relationship processes. *Journal of Personality*, doi:10.1111/jopy.12300
- Atherton BD, Nevels RM, Moore MT. Predicting symptoms of depression from social anhedonia and emotion regulation. *J Nerv Ment Dis* (2015) 203:170–4. doi: 10.1097/NMD.0000000000000262
- Barch, D. M., & Dowd, E. C. (2010). Goal representations and motivational drive in schizophrenia: the role of prefrontal–striatal interactions. *Schizophrenia bulletin*, 36(5), 919-934.
- Barch, D. M., Treadway, M. T., & Schoen, N. (2014). Effort, anhedonia, and function in schizophrenia: reduced effort allocation predicts amotivation and functional impairment. *Journal of abnormal psychology*, 123(2), 387.
- Barch, D. M., Pagliaccio, D., & Luking, K. (2015). Mechanisms underlying motivational deficits in psychopathology: similarities and differences in depression and schizophrenia. In *Behavioral neuroscience of motivation* (pp. 411-449). Springer, Cham.
- Baslet, G., Termini, L., & Herbener, E. (2009). Deficits in emotional awareness in schizophrenia and their relationship with other measures of functioning. *The Journal of nervous and mental disease*, 197(9), 655.
- Barkus, E. & Badcock, J.C. (2019). A transdiagnostic perspective on social anhedonia. *Frontiers in Psychiatry*, 10, e216.
- Bedwell, J.S., Gooding, D.C., Chan, C.C., & Trachik, B.J. (2014). Anhedonia in the age of RDoC. *Schizophrenia Research*, 160, 226-227
- Berlin, I., Givry-Steiner, L., Lecrubier, Y., & Puech, A. (1998). Measures of anhedonia and hedonic responses to sucrose in depressive and schizophrenic patients in comparison with healthy subjects. *European psychiatry*, 13(6), 303-309.
- Berenbaum H, Oltmanns TF (1992) Emotional experience and expression in schizophrenia and depression. *J Abnorm Psychol* 101(1):37–44

- Bodapati, A., Jenkins, L., Sharma, R., & Rosen, C. (2019). Visual memory uniquely predicts anhedonia in schizophrenia but not bipolar disorder. *Journal of Neuropsychology*, 13(1), 136-146.
- Bowie CR, Reichenberg A et al (2006) Determinants of real-world functional performance in schizophrenia subjects: correlations with cognition, functional capacity, and symptoms. *Am J Psychiatry* 163(3):418–425
- Bress JN, Smith E et al (2012) Neural response to reward and depressive symptoms in late childhood to early adolescence. *Biol Psychol* 89(1):156–162
- Blanchard, J.J., Muesser, K.T., Bellack, A.S. (1998). Anhedonia, positive and negative affect, and social functioning in schizophrenia. *Schizophrenia Bulletin*, 24, 413-424
- Blanchard, J.J., Horan, W.P., & Brown, S.A. (2001). Diagnostic differences in social anhedonia: A longitudinal study of schizophrenia and major depressive disorder. *Journal of Abnormal Psychology*, 111, 363-371.
- Blanchard, J.J., Collins, L.M., Aghevli, M., Leung, W.W., & Cohen, A.S. (2011). Social anhedonia and schizotypy in a community sample: The Maryland Longitudinal Study of Schizotypy. *Schizophrenia Bulletin*, 37, 587-602.
- Blanchard, J.J., Park, S.G., Catalano, L.T., & Bennett, M.E. (2015). Social affiliation and negative symptoms in schizophrenia: examining the role of behavioral skills and subjective responding. *Schizophrenia Research*, 168, 491-497.
- Burbridge, J., & Barch, D. (2007). Anhedonia and the Experience of Emotion in Individuals With Schizophrenia. *Journal of Abnormal Psychology*, 116(1), 30-42.
- Burgdörfer, G., & Hautzinger, M. (1987). Psychological and social anhedonia. Evaluation of a research instrument for measuring a basic psychopathologic disorder. *European archives of psychiatry and neurological sciences*, 236(4), 223-229.
- Campellone, T. R., & Kring, A. M. (2018). Anticipated pleasure for positive and negative social interaction outcomes in schizophrenia. *Psychiatry research*, 259, 203-209.

- Campellone, T. R., Truong, B., Gard, D., & Schlosser, D. A. (2018). Social motivation in people with recent-onset schizophrenia spectrum disorders. *Journal of psychiatric research*, *99*, 96-103.
- Chan, R. C., Shi, H. S., Geng, F. L., Liu, W. H., Yan, C., Wang, Y., & Gooding, D. C. (2015). The Chapman psychosis-proneness scales: consistency across culture and time. *Psychiatry research*, *228*(1), 143-149.
- Chapman, L.J., Chapman, J.P., & Raulin, M.L. (1976). Scales for physical and social anhedonia. *Journal of Abnormal Psychology*, *85*, 374-382.
- Chapman, L. J., Chapman, J. P., Kwapil, T. R., Eckblad, M., & Zinser, M. C. (1994). Putatively psychosis-prone subjects 10 years later. *Journal of abnormal psychology*, *103*(2), 17
- Choi, S. H., Lee, H., Ku, J., Yoon, K. J., & Kim, J. J. (2014). Neural basis of anhedonia as a failure to predict pleasantness in schizophrenia. *The World Journal of Biological Psychiatry*, *15*(7), 525-533. Chicago
- Chuang, J. Y., Murray, G. K., Metastasio, A., Segarra, N., Tait, R., Spencer, J., ... & Suckling, J. (2014). Brain structural signatures of negative symptoms in depression and schizophrenia. *Frontiers in psychiatry*, *5*, 116.
- Cicero, D., Martin, E., Becker, T., & Kerns, J. (2014). Reinforcement learning deficits in people with schizophrenia persist after extended trials. *Psychiatry Research*, *220*(3), 760-764.
- Cicero, D. C., Krieg, A., Becker, T. M., & Kerns, J. G. (2016). Evidence for the discriminant validity of the Revised Social Anhedonia Scale from social anxiety. *Assessment*, *23*(5), 544-556.
- Cohen, A.S. & Minor, K.S. (2010). Emotional experience in patients with schizophrenia revisited: meta-analysis of lab studies. *Schizophrenia Bulletin*, *36*, 143-150.
- Culbreth, A. J., Gold, J. M., Cools, R., & Barch, D. M. (2016). Impaired activation in cognitive control regions predicts reversal learning in schizophrenia. *Schizophrenia bulletin*, *42*(2), 484-493.

- Dodell-Feder, D., & Germine, L. (2018). Epidemiological dimensions of social anhedonia. *Clinical Psychological Science*, 6(5), 735-743.
- Dowd, E. C., & Barch, D. M. (2012). Pavlovian reward prediction and receipt in schizophrenia: relationship to anhedonia. *PloS one*, 7(5)
- Downar J, Geraci J et al (2014) Anhedonia and reward-circuit connectivity distinguish nonresponders from responders to dorsomedial prefrontal repetitive transcranial magnetic stimulation in major depression. *Biol Psychiatry* 76(3):176–185
- Durand, D., Strassnig, M., Sabbag, S., Gould, F., Twamley, E. W., Patterson, T. L., & Harvey, P. D. (2015). Factors influencing self-assessment of cognition and functioning in schizophrenia: implications for treatment studies. *European Neuropsychopharmacology*, 25(2), 185-191.
- Edwards, C. J., Cella, M., Tarrier, N., & Wykes, T. (2015). Investigating the empirical support for therapeutic targets proposed by the temporal experience of pleasure model in schizophrenia: A systematic review. *Schizophrenia Research*, 168(1), 120–144.
- Eckblad, M. L., Chapman, L. J., Chapman, J. P., & Mishlove, M. (1982). The revised social anhedonia scale. *Unpublished test*.
- Engel, M., Fritzsche, A., & Lincoln, T. M. (2016). Anticipation and experience of emotions in patients with schizophrenia and negative symptoms. An experimental study in a social context. *Schizophrenia research*, 170(1), 191-197.
- Fawcett, J., Clark, D. C., Scheftner, W. A., & Hedeker, D. (1983). Differences between anhedonic and normally hedonic depressive states. *The American journal of psychiatry*.
- Fortunati, R., Ossola, P., Camerlengo, A., Bettini, E., De Panfilis, C., Tonna, M., ... & Marchesi, C. (2015). Anhedonia in schizophrenia: The role of subjective experiences. *Comprehensive psychiatry*, 62, 152-160.
- Foti D, Hajcak G (2009) Depression and reduced sensitivity to non-rewards versus rewards: evidence from event-related potentials. *Biol Psychol* 81(1):1–8

- Frost, K. H., & Strauss, G. P. (2016). A review of anticipatory pleasure in schizophrenia. *Current Behavioral Neuroscience Reports*, 3(3), 232–247. <http://dx.doi.org/10.1007/s40473-016-0082-5>.
- Fulford, D., Campellone, T., & Gard, D. E. (2018). Social motivation in schizophrenia: How research on basic reward processes informs and limits our understanding. *Clinical psychology review*, 63, 12-24.
- Gard, D. E., Kring, A. M., Gard, M. G., Horan, W. P., & Green, M. F. (2007). Anhedonia in schizophrenia: distinctions between anticipatory and consummatory pleasure. *Schizophrenia research*, 93(1-3), 253-260.
- Germine, L. T., Garrido, L., Bruce, L., & Hooker, C. (2011). Social anhedonia is associated with neural abnormalities during face emotion processing. *Neuroimage*, 58(3), 935-945.
- Gorwood P. Neurobiological mechanisms of anhedonia. *Dialogues Clin Neurosci* (2008), 10:29199.
- Gotlib IH, Hamilton JP et al (2010) Neural processing of reward and loss in girls at risk for major depression. *Arch Gen Psychiatry* 67(4):380–387
- Gooding, D. C., & Pflum, M. J. (2014a). The assessment of interpersonal pleasure: Introduction of the anticipatory and consummatory interpersonal pleasure scale (ACIPS) and preliminary findings. *Psychiatry Research*, 215(1), 237-243.
- Gooding, D. C., & Pflum, M. J. (2014b). Further validation of the ACIPS as a measure of social hedonic response. *Psychiatry Research*, 215, 771-777.
- Gould, F., McGuire, L. S., Durand, D., Sabbag, S., Larrauri, C., Patterson, T. L., ... & Harvey, P. D. (2015). Self-assessment in schizophrenia: Accuracy of evaluation of cognition and everyday functioning. *Neuropsychology*, 29(5), 675.
- Gradin, V. B., Kumar, P., Waiter, G., Ahearn, T., Stickle, C., Milders, M., ... Steele, J. D. (2011). Expected value and prediction error abnormalities in depression and schizophrenia. *Brain*, 134(6), 1751–1764.

- Granholm, E., Ben-Zeev, D., & Link, P. C. (2009). Social disinterest attitudes and group cognitive-behavioral social skills training for functional disability in schizophrenia. *Schizophrenia Bulletin*, 35(5), 874–883. <http://dx.doi.org/10.1093/schbul/sbp072>.
- Green, M.F., Horan, W.P., & Lee, J. (2015). Social cognition in schizophrenia. *Nature Reviews*, 16, 620-631.
- Grimm O, Vollstadt-Klein S et al (2012) Reduced striatal activation during reward anticipation due to appetite-provoking cues in chronic schizophrenia: a fMRI study. *Schizophr Res* 134(2–3):151–157
- Gross, G. M., Mellin, J., Silvia, P. J., Barrantes-Vidal, N., & Kwapil, T. R. (2014). Comparing the factor structure of the Wisconsin Schizotypy Scales and the Schizotypal Personality Questionnaire. *Personality Disorders: Theory, Research, and Treatment*, 5(4), 397.
- Gruzelier, J., & Davis, S. (1995). Social and physical anhedonia in relation to cerebral laterality and electrodermal habituation in unmedicated psychotic patients. *Psychiatry research*, 56(2), 163-172.
- Hallford, D. J., & Sharma, M. K. (2019). Anticipatory pleasure for future experiences in schizophrenia spectrum disorders and major depression: A systematic review and meta-analysis. *British Journal of Clinical Psychology*, 58(4), 357-383.
- Harrow, M., Grossman, L. S., Jobe, T. H., & Herbener, E. S. (2005). Do patients with schizophrenia ever show periods of recovery? A 15-year multi-follow-up study. *Schizophrenia Bulletin*, 31(3), 723-734.
- Herbener, E. S., & Harrow, M. (2002). The course of anhedonia during 10 years of schizophrenic illness. *Journal of Abnormal Psychology*, 111(2), 237.
- Herbener, E. S., Rosen, C., Khine, T., & Sweeney, J. A. (2007). Failure of positive but not negative emotional valence to enhance memory in schizophrenia. *Journal of abnormal psychology*, 116(1), 43.

- Horan, W. P., Green, M. F., Kring, A. M., & Nuechterlein, K. H. (2006). Does anhedonia in schizophrenia reflect faulty memory for subjectively experienced emotions?. *Journal of abnormal psychology*, 115(3), 496.
- Horan, W. P., Wynn, J. K., Kring, A. M., Simons, R. F., & Green, M. F. (2010). Electrophysiological correlates of emotional responding in schizophrenia. *Journal of abnormal psychology*, 119(1), 18.
- Insel, T., Cuthbert, B., Garvey, M., Heinssen, R., Pine, D. S., Quinn, K., ... & Wang, P. (2010). Research domain criteria (RDoC): toward a new classification framework for research on mental disorders.
- Juckel G, Friedel E et al (2012) Ventral striatal activation during reward processing in subjects with ultra-high risk for schizophrenia. *Neuropsychobiology* 66(1):50–56
- Kim, S. J., Hong, Y. J., Kim, M. W., Jung, Y. H., Min, S. R., & Kim, J. J. (2019). Inflexible eye fixation pattern in schizophrenia affecting decision-making on daily life. *Psychiatry research*, 274, 414-420.
- Kiwanuka, J. N., Strauss, G. P., McMahon, R. P., & Gold, J. M. (2014). Psychological predictors of functional outcome in people with schizophrenia. *Schizophrenia research*, 157(1-3), 299-304.
- Kollias, C. T., Kontaxakis, V. P., Havaki-Kontaxaki, B. J., Stamouli, S., Margariti, M., & Petridou, E. (2008). Association of physical and social anhedonia with depression in the acute phase of schizophrenia. *Psychopathology*, 41(6), 365-370.
- Kring, A. M., & Moran, E. K. (2008). Emotional response deficits in schizophrenia: insights from affective science. *Schizophrenia bulletin*, 34(5), 819-834.
- Kring, A. M., & Elis, O. (2013). Emotion deficits in people with schizophrenia. *Annual review of clinical psychology*, 9, 409-433.

- Kring, A. M., & Barch, D. M. (2014). The motivation and pleasure dimension of negative symptoms: neural substrates and behavioral outputs. *European Neuropsychopharmacology*, 24(5), 725-736.
- Kuha, A., Suvisaari, J., Perälä, J., Eerola, M., Saarni, S. S., Partonen, T., ... & Tuulio-Henriksson, A. (2011). Associations of anhedonia and cognition in persons with schizophrenia spectrum disorders, their siblings, and controls. *The Journal of nervous and mental disease*, 199(1), 30-37.
- Kwapil, T.R. (1998). Social anhedonia as a predictor of the development of schizophrenia-spectrum disorders. *Journal of Abnormal Psychology*, 107, 558-565.
- Kwapil, T.R., Barrantes-Vidal, N., Silvia, P.J., 2008. The dimensional structure of the Wisconsin Schizotypy Scales: factor identification and construct validity. *Schizophr. Bull.* 34 (3), 444–457. <http://dx.doi.org/10.1093/schbul/sbm098>.
- Kwapil, T.R., Silvia, P.J., Myin-Germeys, I., Anderson, A.J., Coates, S.A., & Brown, L.H. (2009). The social world of the socially anhedonic: Exploring the daily ecology of asociality. *Journal of Research in Personality*, 43, 103-106.
- Lambert, C., Da Silva, S., Ceniti, A. K., Rizvi, S. J., Foussias, G., & Kennedy, S. H. (2018). Anhedonia in depression and schizophrenia: A transdiagnostic challenge. *CNS neuroscience & therapeutics*, 24(7), 615-623.
- Larquet, M., Coricelli, G., Opolczynski, G., & Thibaut, F. (2010). Impaired decision making in schizophrenia and orbitofrontal cortex lesion patients. *Schizophrenia research*, 116(2-3), 266-273.
- Lee, J. S., Chun, J. W., Kang, J. I., Kang, D. I., Park, H. J., & Kim, J. J. (2012). Hippocampus and nucleus accumbens activity during neutral word recognition related to trait physical anhedonia in patients with schizophrenia: an fMRI study. *Psychiatry Research: Neuroimaging*, 203(1), 46-53.

- Lee, J. S., Kim, E. S., Kim, E. J., Kim, J., Kim, E., Lee, S. K., & Kim, J. J. (2016). The relationship between self-referential processing-related brain activity and anhedonia in patients with schizophrenia. *Psychiatry Research: Neuroimaging*, 254, 112-118.
- Li, Z., Yan, C., Lv, Q. Y., Yi, Z. H., Zhang, J. Y., Wang, J. H., ... & Gur, R. C. (2018). Striatal dysfunction in patients with schizophrenia and their unaffected first-degree relatives. *Schizophrenia research*, 195, 215-221.
- Llerena, K., Park, S.G., Couture, S.M., & Blanchard, J.J. (2012). Social anhedonia and affiliation: Examining behavior and subjective reactions within a social interaction. *Psychiatry Research*, 200, 679-686.
- Loas, G., Salinas, E., Guelfi, J. D., & Samuel-Lajeunesse, B. (1992). Physical anhedonia in major depressive disorder. *Journal of Affective Disorders*, 25(2), 139-146.
- Loas, G., Boyer, P., & Legrand, A. (1999). Anhedonia in the deficit syndrome of schizophrenia. *Psychopathology*, 32(4), 207-219.
- Makowski, C. S., Lepage, M., & Harvey, P. O. (2016). Functional neural correlates of social approval in schizophrenia. *Social cognitive and affective neuroscience*, 11(3), 445–457. <https://doi.org/10.1093/scan/nsv125>
- Mann, C. L., Footer, O., Chung, Y. S., Driscoll, L. L., & Barch, D. M. (2013). Spared and impaired aspects of motivated cognitive control in schizophrenia. *Journal of Abnormal Psychology*, 122(3), 745–755. <https://doi-org.ezproxy.bu.edu/10.1037/a0033069>
- Martin, E. A., Becker, T. M., Cicero, D. C., & Kerns, J. G. (2013). Examination of affective and cognitive interference in schizophrenia and relation to symptoms. *Journal of Abnormal Psychology*, 122(3), 733–744. <https://doi-org.ezproxy.bu.edu/10.1037/a0033956>
- Martin, E.A., Cicero, D.C., Bailey, D.H., Karcher, N.R., & Kerns, J.G. (2016). Social anhedonia is not just extreme introversion: Empirical evidence of distinct constructs. *Journal of Personality Disorders*, 30, 451-468

- Martin, E. A., Siegle, G. J., Steinhauer, S. R., & Condray, R. (2019). Timing matters in elaborative processing of positive stimuli: Gamma band reactivity in schizophrenia compared to depression and healthy adults. *Schizophrenia research*, 204, 111-119
- McCabe C, Woffindale C et al (2012) Neural processing of reward and punishment in young people at increased familial risk of depression. *Biol Psychiatry* 72(7):588–594
- McCarthy, J. M., Bradshaw, K. R., Catalano, L. T., Garcia, C. P., Malik, A., Bennett, M. E., & Blanchard, J. J. (2018). Negative symptoms and the formation of social affiliative bonds in schizophrenia. *Schizophrenia Research*, 193, 225–231
- Meehl, P.E. (1962). Schizotaxia, schizotypy, schizophrenia. *American Psychologist*, 17, 827-838.
- Morgan JK, Olino TM et al (2013) Neural response to reward as a predictor of increases in depressive symptoms in adolescence. *Neurobiol Dis* 52:66–74
- Mote, J., & Fulford, D. (2020). Ecological momentary assessment of everyday social experiences of people with schizophrenia: A systematic review. *Schizophrenia research*, 216, 56-68.
- *Naudin, M., Carl, T., Surguladze, S., Guillen, C., Gaillard, P., Belzung, C., El-Hage, W., & Atanasova, B. (2014). Perceptive biases in major depressive episode. *PloS one*, 9(2), e86832. <https://doi.org/10.1371/journal.pone.0086832>
- *Naudin, M., Mondon, K., El-Hage, W., Perriot, E., Boudjarane, M., Desmidt, T., . . . Atanasova, B. (2015). Taste identification used as a potential discriminative test among depression and Alzheimer's disease in elderly: A pilot study. *Psychiatry Research*, 228(2), 228-232.
- Olino TM, McMakin DL et al (2011) I won, but I'm not getting my hopes up: depression moderates the relationship of outcomes and reward anticipation. *Psychiatry Res* 194(3):393–395
- Olino TM, McMakin DL et al (2014) Reduced reward anticipation in youth at high-risk for unipolar depression: a preliminary study. *Dev Cogn Neurosci* 8:55–64

- Olino, T.M. (2016). Future research direction in the positive valence systems: measurement, development, and implications for youth unipolar depression. *Journal of Clinical Child and Adolescent Psychology*, 45, 681-705.
- Olsen, E., Bjorkquist, O., Bodapati, A., Shankman, S., & Herbener, E. (2015). Associations between trait anhedonia and emotional memory deficits in females with schizophrenia versus major depression. *Psychiatry Research*, 230(2), 323-330.
- Park et al., 2015 I.H. Park, J.W. Chun, H.J. Park, M.S. Koo, S. Park, S.H. Kim, J.J. Kim Altered cingulo-striatal function underlies reward drive deficits in schizophrenia *Schizophr. Res.*, 161 (2–3) (2015), pp. 229-236, 10.1016/j.schres.2014.11.005
- Park, I. H., Kim, J., Ku, J., Jang, H. J., Park, S., Kim, C., . . . Kim, S. I. (2009). Characteristics of social anxiety from virtual interpersonal interactions in patients with schizophrenia. *Psychiatry*, 72(1), 79-93. doi:http://dx.doi.org.ezproxy.bu.edu/10.1521/psyc.2009.72.1.79
- Pechtel P, Dutra SJ et al (2013) Blunted reward responsiveness in remitted depression. *J Psychiatric Res* 47(12):1864–1869
- Pelizza, L., & Ferrari, A. (2009). Anhedonia in schizophrenia and major depression: state or trait?. *Annals of General Psychiatry*, 8(1), 22.
- Pizzagalli DA, Iosifescu D et al (2008) Reduced hedonic capacity in major depressive disorder: evidence from a probabilistic reward task. *J Psychiatric Res* 43(1):76–87
- R Core Team* (2021). *R: A language and environment for statistical computing*. *R Foundation for Statistical Computing, Vienna, Austria*.
URL <https://www.R-project.org/>.
- Rey G, Jouvent R, Dubal S. Schizotypy, depression, and anxiety in physical and social anhedonia. *J Clin Psychology* (2009) 65:695–708. doi: 10.1002/jclp.20577
- Robinson, M. D., & Clore, G. L. (2002). Belief and feeling: evidence for an accessibility model of emotional self-report. *Psychological bulletin*, 128(6), 934.

- Ritsner, M. S., Ratner, Y., Mendyk, N., & Gooding, D. C. (2018). The characterization of social anhedonia and its correlates in schizophrenia and schizoaffective patients. *Psychiatry research, 270*, 922-928.
- Sagud, M., Tudor, L., Šimunić, L., Jezernik, D., Madžarac, Z., Jakšić, N., ... & Pivac, N. (2021). Physical and social anhedonia are associated with suicidality in major depression, but not in schizophrenia. *Suicide and Life-Threatening Behavior, 51*(3), 446-454.
- Savitz, J., Hodgkinson, C. A., Martin-Soelch, C., Shen, P. H., SSDczepanik, J., Nugent, A., ... & Drevets, W. C. (2013). The functional DRD3 Ser9Gly polymorphism (rs6280) is pleiotropic, affecting reward as well as movement. *PloS one, 8*(1).
- Schlosser DA, Fisher M et al (2014) Motivational deficits in individuals at-risk for psychosis and across the course of schizophrenia. *Schizophr Res 158*(1–3):52–57
- Schuck, J., Leventhal, D., Rothstein, H., & Irizarry, V. (1984). Physical anhedonia and schizophrenia. *Journal of Abnormal Psychology, 93*, 342-344
- Sczepanik, J. E., Furey, M. L., Nugent, A. C., Henter, I. D., Zarate Jr, C. A., & Lejuez, C. W. (2017). Altered interaction with environmental reinforcers in major depressive disorder: Relationship to anhedonia. *Behaviour research and therapy, 97*, 170-177.
- Shankman, S. A., Nelson, B. D., Harrow, M., & Faull, R. (2010). Does physical anhedonia play a role in depression? A 20-year longitudinal study. *Journal of affective disorders, 120*(1-3), 170-176.
- Sharp C, Kim S et al (2014) Major depression in mothers predicts reduced ventral striatum activation in adolescent female offspring with and without depression. *J Abnorm Psychol 123*(2):298–309
- Sherdell L, Waugh CE et al (2012) Anticipatory pleasure predicts motivation for reward in major depression. *J Abnorm Psychol 121*(1):51–60
- Silk, J. S., Forbes, E. E., Whalen, D. J., Jakubcak, J. L., Thompson, W. K., Ryan, N. D., ... & Dahl, R. E. (2011). Daily emotional dynamics in depressed youth: A cell phone

ecological momentary assessment study. *Journal of experimental child psychology*, 110(2), 241-257.

Simon, J. J., Biller, A., Walther, S., Roesch-Ely, D., Stippich, C., Weisbrod, M., & Kaiser, S. (2010). Neural correlates of reward processing in schizophrenia—relationship to apathy and depression. *Schizophrenia research*, 118(1-3), 154-161.

Simon, J. J., Zimmermann, J., Cordeiro, S. A., Marée, I., Gard, D. E., Friederich, H. C., ... & Kaiser, S. (2018). Psychometric evaluation of the Temporal Experience of Pleasure Scale (TEPS) in a German sample. *Psychiatry research*, 260, 138-143.

Taylor, S. F., Phan, K. L., Britton, J. C., & Liberzon, I. (2005). Neural response to emotional salience in schizophrenia. *Neuropsychopharmacology*, 30(5), 984-995.

Trémeau, F., Antonius, Cacioppo, Ziwich, Jalbrzikowski, Saccence, . . . Javitt. (2009). In support of Bleuler: Objective evidence for increased affective ambivalence in schizophrenia based upon evocative testing. *Schizophrenia Research*, 107(2-3), 223-231.

Trémeau, F., Antonius, D., Nolan, K., Butler, P., & Javitt, D. (2014). Immediate affective motivation is not impaired in schizophrenia. *Schizophrenia Research*, 159(1), 157-163.

Tso, I. F., Grove, T. B., & Taylor, S. F. (2010). Emotional experience predicts social adjustment independent of neurocognition and social cognition in schizophrenia. *Schizophrenia Research*, 122(1-3), 156-163.

Tso, I. F., Grove, T. B., & Taylor, S. F. (2014). Differential hedonic experience and behavioral activation in schizophrenia and bipolar disorder. *Psychiatry research*, 219(3), 470-476.

Uhl B, Kuehner C et al (2015) Altered neural reward and loss processing and prediction error signalling in depression. *Soc Cogn Affect Neuroscience*

Umesh, S., Nizamie, S. H., Goyal, N., Tikka, S., & Bose, S. (2018). Social anhedonia and gamma band abnormalities as a composite/multivariate endophenotype for schizophrenia: a dense array EEG study. *Early intervention in psychiatry*, 12(3), 362-371.

- Ventura J, Helleman GS et al (2009) Symptoms as mediators of the relationship between neurocognition and functional outcome in schizophrenia: a meta-analysis. *Schizophr Res* 113 (2–3):189–199
- Viechtbauer, W., & Cheung, M. W. (2010). Outlier and influence diagnostics for meta-analysis. *Research synthesis methods*, 1(2), 112–125. <https://doi.org/10.1002/jrsm.11>
- Visser, K. F., Chapman, H. C., Ruiz, I., Raugh, I. M., & Strauss, G. P. (2020). A meta-analysis of self-reported anticipatory and consummatory pleasure in the schizophrenia-spectrum. *Journal of Psychiatric Research*, 121, 68-81.
- Vrieze E, Pizzagalli DA et al (2013) Reduced reward learning predicts outcome in major depressive disorder. *Biol Psychiatry* 73(7):639–645
- Waltz, J., Brown, J., Gold, J., Ross, T., Salmeron, B., & Stein, E. (2015). Probing the Dynamic Updating of Value in Schizophrenia Using a Sensory-Specific Satiety Paradigm. *Schizophrenia Bulletin*, 41(5), 1115-1122.
- Waltz, J. A., Kasanova, Z., Ross, T. J., Salmeron, B. J., McMahon, R. P., Gold, J. M., & Stein, E. A. (2013). The roles of reward, default, and executive control networks in set-shifting impairments in schizophrenia. *PLoS one*, 8(2).
- Wang, J., Huang, J., Yang, X-h., Lui, S.S.Y., Cheung, E.F.C., & Chan, R.C.K. (2015). Anhedonia in schizophrenia: deficits in both motivation and hedonic capacity. *Schizophrenia Research*, 168, 465-474
- Wang, Y. Y., Ge, M. H., Zhu, G. H., Jiang, N. Z., Wang, G. Z., Lv, S. X., ... & Cheung, E. F. (2020). Emotion–behavior decoupling in individuals with schizophrenia, bipolar disorder, and major depressive disorder. *Journal of Abnormal Psychology*.
- Whitton, A. E., Treadway, M. T., & Pizzagalli, D. A. (2015). Reward processing dysfunction in major depression, bipolar disorder and schizophrenia. *Current opinion in psychiatry*, 28(1), 7.

- Wynn, J. K., Horan, W. P., Kring, A. M., Simons, R. F., & Green, M. F. (2010). Impaired anticipatory event-related potentials in schizophrenia. *International Journal of Psychophysiology*, 77(2), 141–149.
- Zhang WN, Chang SH et al (2013) The neural correlates of reward-related processing in major depressive disorder: a meta-analysis of functional magnetic resonance imaging studies. *J Affect Disord* 151(2):531–539
- Zhang, B., Lin, P., Shi, H., Öngür, D., Auerbach, R. P., Wang, X., ... & Wang, X. (2016). Mapping anhedonia-specific dysfunction in a transdiagnostic approach: an ALE meta-analysis. *Brain imaging and behavior*, 10(3), 920-939.
- Zou, Y. M., Ni, K., Yang, Z. Y., Li, Y., Cai, X. L., Xie, D. J., ... & Shum, D. H. (2018). Profiling of experiential pleasure, emotional regulation and emotion expression in patients with schizophrenia. *Schizophrenia research*, 195, 396-401

Table 1a.
Study and SSD participant characteristics

Citation	Sample Size		% Male		Mean Age		Dep. Sx. Severity	Neg. Sx. Severity
	SZ	CN	SZ	CN	SZ	CN		
Barch et al. (2008)	66	44	65	66	37.1	36.8	23.65	36.0
Barch et al. (2014)	59	39	57.6	48.7	39.3	37.4	18.10	40.0
Baslet et al. (2009)	21	20	76.2	60	36.86	34.4		
Bererril et al. (2011)	38	32	66	66	36.66	39.19		36.80
Berlin et al. (1998)	20	20	52	35	30	47	15	61.22
Blanchard et al. (1998)	43	15	63	53	36.14	36.4		
Bodapati et al. (2019)	38	43	53	45	39.47	38.26	52.64	
Burbridge et al. (2007)	49	47	63	49	39.4	37.4		
Catalano et al. (2019)	41	29	68.3	69	38	39.1	9.30	35.71
Cohen et al. (2005)	73	22	75	68	41.67	38.95		
Ceaser et al. (2016)	22	20	75	55	40.41	33.65		
Cicero et al. (2014)	54	32	87	91	41.46	43	47.33	40.62
Choi et al. (2014)	15	17	66.7	58.8	29.1	29.1		33.47
Culbreth et al. (2016)	58	40	66.7	52.8	37	36.36	17.67	24.00
Dowd et al. (2012)	29	22	72	70	31.44	33.2		59.00
Fortunati et al. (2015)	53	46	60.4	50	40.1	38.3	9.92	44.65
Gard et al. (2007)	28	29	64.7	50	48.06	46.62		
Gruzelier et al. (1993)	7	11	42.9	54.5				
Herbener et al. (2007)	33	28	57	46.4	37.55	40.82		30.61
Horan et al. (2006)	30	31	83	68	46.2	40.5		
Horan et al. (2010)	38	36	81.6	74.3	44.5	38.5	35.71	46.00
Kim et al. (2019)	23	23	47.8	47.8	33.35	34.04		40.02
Kiwanuka et al. (2014)	100	78	68	70	41.14	40.68		57.20

Note: SSD = Schizophrenia Spectrum Disorder, CN = Control, Prop. Dep. Sx. Severity = Depression Symptom Severity, Neg. Sx. Severity = Negative Symptom Severity

Table 1.b
Study and SSD participant characteristics

Citation	Sample Size		% Male		Mean Age		Dep. Sx. Severity	Neg. Sx. Severity
	SZ	CN	SZ	CN	SZ	CN	SZ	SZ
Kuha et al. (2011)	91	67	51.6	49.2	44.5	47		
Lee et al. (2012)	14	16	64.2	43.8	29.9	30.3		39.18
Lee et al. (2016)	20	25	70	64	43.9	41.5		34.08
Li et al. (2018)	26	26	57.7	57.7	22.77	24.6		32.90
Makowski et al. (2016)	15	15	73.3	60	33.1	35.2	4.074	17.91
Mann et al. (2013)	54	39	59.3	48.7	38.35	36.5		31.60
Martin et al. (2013)	48	28	83.3	92.8	40.78	43.6		
Martin et al. (2019)	15	23	53	47.8	41.47	31.67	21.698	
Olsen et al. (2015)	25	33	0	0	41.56	38.39	16.846	28.25
Park et al. (2015)	20	20	50	40	28.6	26.1		35.10
Park et al. (2009)	27	27	51.9	51.9	28.5	26.5		41.43
Tremeau et al. (2009)	64	32	84.4	81.3	37	37	8.167	9.091
Tremeau et al. (2014)	120	62	80	74	38.1	34.5	6.5	41.63
Tso et al. (2010)	33	33	66.7	69	38.5	38.2		40.48
Tso et al. (2014)	39	36	58.9	63.8	41.5	40.4	18.10	21.63
Umesh et al. (2018)	20	20	100	100	29.8	29.8	18.095	42.86
Waltz et al. (2015)	42	44	76.2	61.4	37.9	38.6		30.00
Waltz et al. (2013)	29	21	82.8	71.46	39.6	39.6		29.00
Wang et al. (2020)	41	43	43.9	44.2	27.3	27.9	25.846	27.10
Wynn et al. (2010)	34	36	78.09	66.7	44.6	38.6		47.60
Zhorinitsky et al. (2012)	23	25	60.9	64	39.9	40.		31.83
Zou et al. (2018)	146	73	56.2	49.3	36.29	36.0	25.302	38.14

Note: SSD = Schizophrenia Spectrum Disorder, CN = Control, Dep. Sx. Severity = Depression Symptom Severity, Neg. Sx. Severity = Negative Symptom Severity

Table 2

Study and MDD participant characteristics

Citation	Sample Size		% Male		Mean Age		Dep. Sx. Severity MDD
	MDD	CN	MDD	CN	MDD	CN	
Berlin et al. (1998)	20	20	35	35	47	47	42
Chuang et al. (2014)	24	20	70	80	33.08	34.3	5.96
Naudin et al. (2014)	22	41	27	41	33.2	34	37.1
Naudin et al. (2015)	20	24	20	38	64.9	67.4	29.12
Savitz et al. (2012)	10	26	20	42.3	38.2	34.5	24.3
Olsen et al. (2015)	18	33	0	0	35.94	38.39	15
Szczepanik et al. (2017)	21	23	61.9	56.5	35.5	31.8	32.7
Wang et al. (2020)	44	43	34	44.1	28.66	27.93	15.11

Note: MDD = Major Depression Disorder, CN = Control, Dep. Sx. Severity = Depression Symptom Severity

Table 3. Results of meta-analyses and meta-regression analyses: SSDs

SSD	Hedge's G	95%CI	Q	I ²	p
Physical Anhedonia	0.89 ***	0.78 1.0	83.9	51.2%	<0.0001
Social Anhedonia	0.93***	0.67 1.19	101.3	56.6%	<0.0001
Moderators (Phys Soc)	β	SE	95%CI Low	95%CI Hi	p
Age	<0.01 -0.01	0.01 0.03	-0.02 -0.07	0.03 0.04	0.72 0.57
%Male	<0.01 0.01	<0.01 0.01	-0.01 -0.01	0.01 0.03	0.74 0.18
%white	-0.01 -0.01	<0.01 <0.01	-0.01 -0.01	0.004 0.01	0.30 0.80
%Atypical AP	-0.01 -0.02	0.004 0.02	-0.01 -0.06	0.002 0.01	0.12 0.20
NegSx	0.01 -0.02	0.01 0.02	-0.01 -0.05	0.02 0.02	0.15 0.39
DepSx	-0.01 0.04	0.01 0.03	-0.02 -0.02	0.01 0.09	0.37 0.21
	Hedge's G	95%CI	Q	I ²	p
Phys. vs. Soc. within-group	-0.50**	-0.83 -0.17	26.0	0.0%	<0.01
Moderators	β	SE	95%CI	p	
Age*	-0.08	0.03	-0.14 -0.02	<0.05	
%Male	<-0.01	0.01	-0.03 0.02	0.66	
%White	0.005	0.02	-0.03 0.04	0.74	
% Atypical AP	<0.01	0.01	-0.02 0.03	0.60	
NegSx	<0.001	0.01	-0.03 0.03	0.97	
DepSx	-0.03	0.02	-0.08 0.01	0.15	

Note: Atypical AP= Atypical Antipsychotic, NegSx= Negative Symptom Severity, DepSx=Depressive Symptom Severity

Table 4. Results of meta-analyses and meta-regression analyses: Major Depressive Disorder

Major Depressive Disorder	Hedge's G	95%CI	Q	I ²	p
Physical Anhedonia	1.15**	0.94 1.37	10.18	31.3%	<0.0001
Social Anhedonia	1.29**	1.07 1.51	11.45	38.9%	<0.0001
Moderators (Phys Soc)	β	SE	95%CI Low	95% Hi	p
Age	-0.00 0.01	0.01 0.01	-0.03 -0.03	0.03 0.02	0.94 0.70
%Male	0.01 0.001	0.01 0.01	-0.00 -0.01	0.02 0.02	0.10 0.94
DepSx	0.02* 0.00	0.01 0.01	0.001 -0.02	0.03 0.02	0.03 0.79
Phys. vs. Soc. within-group	Hedge's G	95%CI	Q	I ²	p
	-0.72	-2.92 -1.47	0.30	0.0%	0.52
Moderators	β	SE	95% CI	p	
Age	-0.003	0.1	-0.20 0.20	0.98	
%Male	0.01	0.06	-0.10 0.13	0.83	
DepSx	0.02	0.07	-0.13 0.16	0.83	

Note: Atypical AP= Atypical Antipsychotic, DepSx=Depression Symptoms Severity




Fig 1 Study Selection Process SSDs




Fig 2 Study Selection Process MDD

A TRANSDIAGNOSTIC REVIEW OF ANHEDONIA

40




Fig 3. Forest Plot of differences in physical anhedonia between SZ & controls




Fig 4 Forest Plot of differences in social anhedonia between SZ & controls. One outlier (Umesh et al.) removed from plot.




Fig 5. Forest Plot of differences between social and physical anhedonia within SZ




Fig 6. Forest Plot of Differences in Physical Anhedonia Between MDD and Controls




Fig 7. Forest plot of differences on Social Anhedonia between MDD and HC

Source	SMD (95% CI)
Berlin et al	0.13 [-5.24; 5.51]
Chuang et al	-0.66 [-9.46; 8.15]
Naudin et al	-0.71 [-6.55; 5.13]
Naudin et al	-0.96 [-6.86; 4.93]
Savitz et al	-1.23 [-8.03; 5.57]
Olsen et al	-1.46 [-6.77; 3.84]
Szczepanik et al	-1.33 [-9.15; 6.49]
Wang et al	0.10 [-5.84; 6.04]
Total	-0.72 [-2.92; 1.47]
95% PI	[-3.46; 2.02]
Heterogeneity: $\chi^2_7 = 0.30$ ($P > .99$), $I^2 = 0\%$	




Fig 8. Forest plot of Differences Between Physical & Social Anhedonia within MDD




Fig 9. Funnel Plot of Differences in Physical Anhedonia Between SSDs and Controls




Fig 10. Funnel Plot of Differences in Social Anhedonia Between SSDs and Controls




Fig 11. Funnel Plot of Differences Between Physical and Social Anhedonia within SSDs




Fig 12. Funnel Plot of Differences in Physical Anhedonia Between MDD and Controls




Fig. 13 Funnel Plot for Differences in Social Anhedonia Between MDD & Controls




Fig. 14 Funnel Plot for Differences between Social and Physical Anhedonia within MDD